

Multiple Scattering And Volume-Roughness Interactions In Sea Bed Acoustics

Anatoliy N. Ivakin
Andreev Acoustics Institute, Shvernika 4, Moscow 117036, Russia
email: aniva@glasnet.ru

Darrell R. Jackson
Applied Physics Laboratory,
College of Ocean and Fishery Sciences,
University of Washington, Seattle, Washington 98105
phone: (206) 543-1359 fax: (206) 543-6785 email: drj@apl.washington.edu
Award #: N00014-97-1-0615

LONG-TERM GOAL

The objective of this research is to improve understanding of bottom acoustic scattering at mid- and high-frequencies and to improve conventional algorithms for remote acoustic characterization of marine sediments.

OBJECTIVES

The primary scientific objective is to develop improved theoretical models for seabed scattering using approximations having a wider range of applicability than those presently available.

APPROACH

The seabed is treated as a multi-layered fluid medium with rough interfaces and volume inhomogeneities covering a rough inhomogeneous elastic halfspace (basement). This makes it possible to unify scattering models for a wide frequency range and for different seabed types. A general description of the scattered field is given in terms of the plane wave scattering amplitude and is based on a unified perturbation approach to volume and roughness scattering [1]. This approach can permit in principle the consideration of multiple scattering in the complex seabed medium with strong volume and roughness irregularities which in general can be correlated. As a first step, the approach gives the first-order perturbation solution in a unified form more convenient for describing both volume and roughness components of scattering. This provides direct relationships between characteristics of the scattered field and parameters of the seafloor. Average seabed parameters can be arbitrary functions of depth, and statistical properties of the seabed are described by power-law spectra of seabed roughness and the sediment volume inhomogeneity in forms consistent with available data.

WORK COMPLETED

A first-order model of seabed scattering has been developed in detail based on realistic assumptions about sediment properties, including roughness and inhomogeneity, vertical stratification (gradients and/or layering), and ability to support both compressional and shear waves [1-3]. Corresponding codes have been developed and numerical examples for various seabed types and particular models of scattering have been presented and analyzed in recent publications [4-9].

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 1998		2. REPORT TYPE		3. DATES COVERED 00-00-1998 to 00-00-1998	
4. TITLE AND SUBTITLE Multiple Scattering and Volume-Roughness Interactions in Sea Bed Acoustics				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Washington, Applied Physics Laboratory, Seattle, WA, 98105				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM002252.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

RESULTS

The model and codes developed have made it possible to analyze the sensitivity of measurable scattering characteristics (e.g., the scattering strength and its angular-frequency dependencies) with respect to various seabed parameters and determine those parameters that can be considered as classification clues for remote acoustic characterization of marine sediments. Continued model-data comparisons using data from the CBBL experiment at Key West have confirmed the earlier conclusion that gradient effects are not responsible for slight oscillations observed in the bistatic scattering strength angular dependence. Examples of strong gradient effects on seabed volume and roughness scattering are given, e.g., in [7].

IMPACT/APPLICATION

The models of seabed scattering developed in this research will provide a better understanding of bottom acoustic interaction at mid- and high-frequencies and can be used as a basis for improved algorithms for remote acoustic inversions for seafloor properties.

TRANSITIONS

The results of this work are being adapted in practical models for seabed scattering. For example, a high-frequency bistatic scattering model funded by the ONR Torpedo Environments Program (6.2) incorporates the elastic scattering model developed as part of this work.

RELATED PROJECTS

This research is conducted jointly with the separately funded work of D.R. Jackson and comparisons with CBBL data have been carried out in collaboration with Kevin Williams. The approaches and models developed in this research are relevant to acoustic penetration and multiple scattering issues arising within the ONR Departmental Research Initiative on high-frequency sound interaction with the seafloor.

REFERENCES

1. Ivakin, A.N., 1998. "A unified approach to volume and roughness scattering," J. Acoust. Soc. Am., 103, 827-837.
2. Ivakin, A.N., 1997, "A unified model for seabed volume and roughness scattering," in High Frequency Acoustics in Shallow Water, N.G.Pace, E.Pouliquen, O.Bergem and A.P.Lyons, eds., SACLANT, La Spezia, Italy, pp.267-273.
3. Jackson D.R. and A.N. Ivakin, 1998, "Scattering from elastic sea beds: First-order theory," J. Acoust. Soc. Am., 103, 336-345.
4. Ivakin, A.N. and D.R. Jackson, 1998, "Effects of shear elasticity on sea bedscattering: Numerical examples," J. Acoust. Soc. Am., 103, 346-354.

5. A.N. Ivakin, 1998, "Models of volume and roughness scattering in stratified seabeds ", in Proc. 16th International Congress on Acoustics and 135th Meeting of Acous. Soc. Amer., Seattle, WA, 4, 2698-2699.
6. D.R. Jackson and A.N. Ivakin, 1998, "A practical model for high-frequency seabed bistatic scattering strength ", in Proc. 16th Internat. Congress on Acoustics and 135th Meeting of Acous. Soc. Amer., Seattle, WA, 4, 2693-2694.
7. A.N. Ivakin, 1998, "Models of seafloor roughness and volume scattering", in Oceans'98 conference Proceedings, 28 Sept.-1 Oct. 1998, Nice, 1, 518-521.
8. A.N. Ivakin, 1997, "First-order model for bottom volume and roughness scattering", in Shallow-Water Acoustics, R.H.Zhang and J.X.Zhou, eds., China Ocean Press, Beijing, pp. 359-364.
9. A.N. Ivakin, 1997, "Effects of cross correlation between different types of medium perturbations on seabed scattering", J. Acous. Soc. Amer., 102, Pt.2, 216.